

The Araucaria Project. The Distance to the Sculptor Group Galaxy NGC 55 from a Newly Discovered Abundant Cepheid Population.¹

Grzegorz Pietrzyński

Universidad de Concepción, Departamento de Fisica, Astronomy Group, Casilla 160-C,
Concepción, Chile

Warsaw University Observatory, Al. Ujazdowskie 4, 00-478, Warsaw, Poland

Wolfgang Gieren

Universidad de Concepción, Departamento de Fisica, Astronomy Group, Casilla 160-C,
Concepción, Chile

Igor Soszyński

Universidad de Concepción, Departamento de Fisica, Astronomy Group, Casilla 160-C,
Concepción, Chile

Warsaw University Observatory, Aleje Ujazdowskie 4, PL-00-478, Warsaw, Poland

Andrzej Udalski

Warsaw University Observatory, Aleje Ujazdowskie 4, PL-00-478, Warsaw, Poland

Fabio Bresolin

Institute for Astronomy, University of Hawaii at Manoa, 2680 Woodlawn Drive, Honolulu
HI 96822, USA

Rolf-Peter Kudritzki

Institute for Astronomy, University of Hawaii at Manoa, 2680 Woodlawn Drive, Honolulu
HI 96822, USA

Ronald Mennickent

Universidad de Concepción, Departamento de Fisica, Astronomy Group, Casilla 160-C,
Concepción, Chile

Alistair Walker

Cerro Tololo Inter-American Observatory, Casilla 603, La Serena, Chile

Alejandro Garcia

¹Based on observations obtained with the 1.3 m Warsaw telescope at Las Campanas Observatory, Chile.

Universidad de Concepción, Departamento de Física, Astronomy Group, Casilla 160-C,
Concepción, Chile

Olaf Szewczyk
Warsaw University Observatory, Aleje Ujazdowskie 4, PL-00-478, Warsaw, Poland

Michał Szymański
Warsaw University Observatory, Aleje Ujazdowskie 4, PL-00-478, Warsaw, Poland

Marcin Kubiak
Warsaw University Observatory, Aleje Ujazdowskie 4, PL-00-478, Warsaw, Poland

Łukasz Wyrzykowski
Warsaw University Observatory, Aleje Ujazdowskie 4, PL-00-478, Warsaw, Poland

ABSTRACT

We have detected, for the first time, Cepheid variables in the Sculptor Group SB(s)m galaxy NGC 55. From wide-field images obtained in the optical V and I bands during 77 nights in 2002-2003, we have found 143 Cepheids with periods ranging from 5.6 to 175.9 days. 133 of these objects have periods longer than 10 days, making NGC 55 to-date the galaxy with the largest known number of long-period Cepheids in the Sculptor Group. We construct period-luminosity relations from our data and obtain distance moduli corrected for the small foreground reddening to NGC 55 of 26.79 ± 0.04 mag (internal error) in V, 26.66 ± 0.03 mag in I and 26.40 ± 0.05 mag in the reddening-independent V-I Wesenheit index. The trend of increasing distance moduli with shorter wavelength hints at the existence of significant reddening intrinsic to NGC 55 which affects the measured Cepheid magnitudes. From our data, we determine the intrinsic mean reddening of the Cepheids in NGC 55 as $E(B-V) = 0.102$ mag which brings the distance determinations from the different bands into excellent agreement. Our best distance estimate for NGC 55 from the present optical Cepheid photometry is 26.40 mag ± 0.05 mag (internal error) ± 0.09 mag (systematic error). This value is tied to an assumed LMC distance of 18.50 mag. Our quoted systematic error of the present NGC 55 Cepheid distance does not take into account the current uncertainty on the distance of the fiducial LMC galaxy itself.

Within the small respective uncertainties, the Sculptor Group galaxies NGC 55 and NGC 300 are at the same distance of 1.9 Mpc, strengthening the case for a physical association of these galaxies.

Subject headings: distance scale - galaxies: distances and redshifts - galaxies:
individual: NGC 55 - galaxies: stellar content - stars: Cepheids

1. Introduction

In our ongoing Araucaria Project, we are improving the accuracy of several of the most important stellar distance indicators with the aim to measure the distances to nearby galaxies, out to about 10 Mpc, with an accuracy of 3 percent with respect to the Large Magellanic Cloud (LMC). The feasibility to reach this accuracy using a combination of optical and near-infrared photometry of *Cepheid variables* has already been demonstrated for the Local Group irregular galaxies IC 1613 (Pietrzyński et al. 2006a), NGC 6822 (Gieren et al. 2006; Pietrzyński et al. 2004) and NGC 3109 (Pietrzyński et al. 2006b, Soszyński et al. 2006). Our list of targets also includes four of the five principal late-type galaxies in the loose neighboring group of galaxies in Sculptor. For NGC 300, a near face-on spiral member of the Sculptor Group at a distance of 1.88 Mpc, we were able to achieve the same 3 percent accuracy for its distance determination from the Cepheid approach (Gieren et al. 2005a). For all these galaxies so far studied in the Araucaria Project, we found clear evidence for the existence of substantial intrinsic dust absorption, in addition to the foreground absorption produced inside the Milky Way. This result underlines the importance to use infrared data to obtain the most reliable distance estimates, eliminating intrinsic reddening as the dominating source of systematic error in optical photometric studies. Our previous studies also show that in the case of Cepheids, the reddening-independent *Wesenheit index* is able to provide a distance result which is much less affected by reddening than the distance estimates in the V and I bands, provided excellent optical light curves of the variables are available. A Cepheid distance derived from the (V-I) Wesenheit index (for a definition see our previous papers cited above) is therefore a good approximation to the more accurate distance which can be obtained from near-infrared data, and allows an important consistency check on the distance result for a given galaxy from infrared photometry of its Cepheid variables.

NGC 55, our target galaxy studied in this paper, is a highly inclined late-type galaxy classified as SB(s)m in the NASA Extragalactic Database. Although its high inclination makes it difficult to study its morphology, NGC 55 resembles to some extent the LMC but is more massive. From color images as the one shown in Gieren et al. (2005b), it is evident that NGC 55 contains an abundant population of young blue stars and should therefore contain a sizable population of Cepheids, which have so far not been searched for in this

galaxy. The few previous distance determinations which have been attempted for NGC 55 have yielded an unsatisfactorily large range, from 1.34 Mpc from carbon stars (Pritchett et al. 1987) to 2.3 Mpc from planetary nebulae (Van de Steene et al. 2006). In between falls the determination of 1.8 Mpc obtained by Karachentsev et al. (2003) who used the Tully-Fisher relation, tied to distances of other dwarf galaxies in the Sculptor Group measured with the tip of the red giant branch method. All these distance determinations are likely to be plagued by rather large systematic uncertainties. For instance, the recent determination of Van de Steene et al. (2006) from the PNLf method depends rather sensitively on the assumed oxygen abundance of the planetary nebulae in NGC 55 used in the construction of the luminosity function, which is quite uncertain at the present time (e.g. Tuellmann et al. 2003). The detection of a sizable population of Cepheid variables in NGC 55 has therefore a strong potential to improve the situation and provide a more accurate distance determination for this galaxy which we need in the context of the principal science goal of the Araucaria Project, viz. to determine the effect of metallicity on the physical parameters used as a distance indicator for the different stellar standard candles we study. We remark here that we are obtaining, in a parallel effort, chemical abundances for oxygen and some other elements from the spectra of a large number of blue supergiant stars in NGC 55 which we have obtained with the ESO VLT. These abundances will not only provide a mean metallicity for NGC 55 but also allow to determine the abundance *gradient* in its disk if existent, as we were able to do in the case of NGC 300 (Urbaneja et al. 2005).

The purpose of this paper is to present the results of an extensive wide-field imaging survey for Cepheid variables in NGC 55 which we conducted in 2002 and 2003 at the Warsaw 1.3-m telescope at Las Campanas Observatory. As we will show in the following sections, we have discovered a surprisingly large population of long-period Cepheids in this galaxy. We will describe our observations, data reductions and calibrations in section 2, and will present the catalog of photometric properties of the Cepheid variables in NGC 55 in section 3. In section 4 we will determine the period-luminosity relations in the V, I and Wesenheit bands from our data and provide a new distance determination to NGC 55. We will discuss our results in section 5, and summarize our conclusions in section 6.

2. Observations, Reductions and Calibrations

All the data presented in this paper were collected with the Warsaw 1.3-m telescope at Las Campanas Observatory. The telescope was equipped with a mosaic $8k \times 8k$ detector, with a field of view of about 35×35 arcmin and a scale of about 0.25 arcsec/pix. For more instrumental details on this camera, the reader is referred to the OGLE website:

<http://ogle.astrouw.edu.pl/~ogle/index.html>. V and I band images of NGC 55 were secured during 77 different nights. The exposure time was set to 900 seconds in both filters. Most of the observations were obtained in 2002 between July 3 and November 21. Additional observations were made during three photometric nights in May 2003 in order to accurately calibrate the data and improve the periods of the detected variable stars.

Preliminary reductions (i.e. debiasing and flatfielding) were done with the IRAF² package. Then, the PSF photometry was obtained for all stars in the same manner as in Pietrzyński, Gieren and Udalski (2002). Independently, the data were reduced with the OGLE III pipeline based on the image subtraction technique (Udalski 2003; Woźniak 2000).

In order to calibrate our photometry onto the standard system our target was monitored during three photometric nights (2003 May 11, 17 and 28) together with some 25 standard stars from the Landolt fields spanning a wide range of colors ($-0.14 < V-I < 1.43$), and observed at very different airmasses. Since in principle the transformation equations for each of the eight chips may have different color coefficients and zero points, the selected sample of standard stars was observed on each of the individual chips, and transformation coefficients were derived independently for each chip, on each night. Comparison of the photometry obtained on the different nights revealed that the internal accuracy of zero points in both V and I bands is better than 0.02 mag.

To correct the variation of the zero points in V and I over the mosaic, the "correction maps" established by Pietrzyński et al. (2004) were used. These maps were already applied to correct photometry obtained in the field of NGC 6822 (Pietrzyński et al. 2004) and NGC 3109 (Pietrzyński et al. 2006b) with the same camera. Comparison with other studies revealed that these maps allow to correct for the zero point variations across the mosaic field in a way that the residuals do not exceed 0.02-0.03 mag.

Two fields in the SMC which contain a huge number of uniformly distributed stars with very accurate photometry established by the OGLE team (Udalski et al. 1999) were monitored on all three nights together with our target and standard fields, and were calibrated in the same manner as the NGC 55 observations. Comparison of OGLE II and our photometry shows that on average the difference in the zero points between these two photometric sets is smaller than 0.02 mag on each chip.

Alcaino and Liller (1983) established a sequence of secondary standards in the field of NGC 55. In our database 7 stars were found to be common to this sequence (e.g. the

²IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the NSF.

remaining ones were either saturated or located outside our field of view). The comparison of our V band photometry for these objects with that of Alcaïno and Liller, which is presented in Table 1, demonstrates that the zero points of the two datasets agree to within 0.02 mag.

Summarizing the previous discussion, we may conclude that all internal and external checks we made indicate that the zero points of our data are accurate to 0.02 mag in both V and I, and that their possible variations over the mosaic fields are smaller than 0.02 mag.

3. The Catalog of Cepheids

All stars with photometry in NGC 55 were searched for photometric variations with periods between 0.2 and 200 days, using the analysis of variance algorithm (Schwarzenberg-Czerny 1989). In order to distinguish Cepheids from other types of variable stars, we used the same criteria given in our initial paper reporting on the discovery of Cepheids in NGC 300 (Pietrzyński et al. 2002). The light curves of all Cepheid candidates were approximated by Fourier series of order 4. We then rejected those objects with V amplitudes smaller than 0.4 mag, the approximate lower limit amplitude for classical Cepheids. This procedure also helps in screening our Cepheid sample from the inclusion of strongly blended objects. For the variable stars passing our selection criteria, mean V and I magnitudes were derived by integrating their light curves which had been previously converted onto an intensity scale, and converting the results back to the magnitude scale. Because of the very good quality and phase coverage of the light curves, the statistical accuracy of the derived mean magnitudes was typically 0.01 mag for the brighter variables, increasing to about 0.03 mag for the faintest Cepheids in our catalog. We ended up with a list of 143 bona fide classical Cepheids with periods between 175.9 and 5.6 days. The number of long-period Cepheids in NGC 55 is surprisingly large, with 133 of the detected Cepheids having periods longer than 10 days. This makes NGC 55 even richer in its long-period Cepheid population than NGC 300, the other Sculptor galaxy we had previously surveyed for Cepheids (Gieren et al. 2004). There is certainly a note of caution as to the true Cepheid nature of the 3 objects with periods in excess of 100 days; the time baseline of our photometry is possibly too short to establish their Cepheid character beyond any doubt. Future observations will clarify this question. For this reason, we will exclude these variables in our distance determination in the next section, as we did in the case of the Local Group galaxy NGC 6822 in which we had also found a number of very long-period Cepheid-like variables (Gieren et al. 2006; Mennickent et al. 2006).

Table 2 gives the individual photometric V and I observations for all Cepheid variables

we found in NGC 55. The full Table 2 is available in electronic form. In Table 3, we present their identifications, equatorial coordinates, periods, times of maximum brightness in V, and their mean intensity magnitudes in the V, I and W bands. The accuracy of the periods is typically about $10^{-3} * P$ days, good enough to be no significant source of error in the construction of the period-luminosity relations discussed in the next section.

In Fig. 1, we show a mosaic image of NGC 55 from our data with the positions of the detected Cepheids overplotted. Fig. 2 presents light curves for some of the Cepheids which are representative in quality for other variables in our catalog of similar periods, showing that the mean magnitudes of the Cepheids can indeed be derived quite accurately from our data even for the shortest-period and faintest Cepheids in our sample. In Fig. 3 we show the locations of the Cepheids in the V, V-I color-magnitude diagram for NGC 55 which we have constructed from our data. The diagram shows that the Cepheids indeed do fill the expected zone in the CMD delineating the Cepheid instability strip, presenting supporting evidence that our classification of the objects in Table 3 as Cepheids seems correct in all cases.

4. Period-Luminosity Relations and Distance Determination

Although a more accurate distance to NGC 55 will be derived once near-infrared follow-up photometry will become available for a substantial subsample of the Cepheids in our catalog, we can already clearly improve on the existing distance determinations for NGC 55 with the optical Cepheid photometric data we have provided in this paper. In Figs. 4-6, we show the period-luminosity (PL) diagrams obtained from the data in our Cepheid catalog for the 130 stars with periods in the range from 10-100 days. PL relations are plotted for the V, I and W magnitudes, where the (V-I) Wesenheit magnitude is defined in its usual way (see previous papers in the Araucaria Project). The PL relation in V is well defined although with somewhat larger scatter than in the case of NGC 300 at a similar distance (Pietrzyński et al. 2002; Gieren et al. 2004). This is likely due to an increased size of differential reddening in NGC 55 due to its high inclination with respect to the line of sight, as compared to NGC 300 which is oriented almost face-on. As expected if differential reddening indeed causes much of the scatter in Fig. 4, the I-band PL relation in Fig. 5 has a clearly smaller dispersion. While it should be expected that the W-band PL relation has an even smaller dispersion due to its expected insensitivity to reddening, Fig. 6 shows that this is not the case and that the dispersion is instead rather similar to the one in the V-band PL relation. This is probably because the photometric accuracy of the data especially for the shorter-period Cepheids is not quite as high as needed for the

W-band technique to show its full power. It is also possible that crowding and blending of the Cepheids with bright unresolved companion stars in this high-inclination galaxy is a more significant problem than in the other galaxies we have studied so far, invalidating the power of the Wesenheit approach to some extent. This has to be kept in mind when discussing the accuracy of our present distance determination.

In order to obtain the distance to NGC 55, we have carried out least-squares fits to a line to the various PL diagrams. These free fits yield the following slopes for the PL relations: -2.31 ± 0.18 , -2.56 ± 0.12 and -2.85 ± 0.20 in V, I and W_I , respectively. These slope values are somewhat shallower than the corresponding slopes for the LMC Cepheid sample studied by the OGLE Project, but they are still consistent with the OGLE LMC PL relation slopes of -2.775 in V, -2.977 in I and -3.300 in W_I (Udalski 2000) at the level of 2σ . We therefore keep our standard procedure adopted in our previous papers in this series to use the OGLE LMC slopes to derive the distance of our target galaxies relative to the LMC. Forcing the slopes to the respective OGLE values leads to the following equations for the NGC 55 Cepheid PL relations:

$$V = -2.775 \log P + (25.401 \pm 0.038) \text{ mag}$$

$$I = -2.977 \log P + (24.778 \pm 0.026) \text{ mag}$$

$$W_I = -3.300 \log P + (23.765 \pm 0.045) \text{ mag}$$

Adopting 18.50 mag for the LMC distance modulus as we did in our previous papers, and using $E(B-V) = 0.014$ mag as the foreground reddening towards NGC 55 (Schlegel et al. 1998) together with the reddening law of the same authors ($A_V = 3.24 E(B-V)$, $A_I = 1.96 E(B-V)$), we obtain the following distance moduli for NGC 55 from the three different bands:

$$(m - M)_0 (W_I) = 26.40 \pm 0.05 \text{ mag}$$

$$(m - M)_0 (I) = 26.66 \pm 0.03 \text{ mag}$$

$$(m - M)_0 (V) = 26.79 \pm 0.04 \text{ mag}$$

These values show the same tendency of increasing distance moduli towards the bands with the higher reddening sensitivity that we have seen for the other galaxies previously studied in this series, suggesting the existence of significant internal reddening in NGC 55 in addition to the foreground reddening produced in the Milky Way.

This is of course no surprise because high-resolution images of NGC 55 clearly reveal the existence of significant dust absorption in the regions where the young, blue stars are concentrated. Since the difference of the distance moduli in V and I above is identical to the reddening $E(V-I)$, we obtain from our data $E(V-I) = 26.79 - 26.66 = 0.13$ mag. With $E(B-V) = E(V-I)/1.28$, this translates in a (B-V) color excess of $E(B-V) = 0.102$ mag, which then leads to an extinction-corrected true distance modulus of $(m - M)_0 = 26.46$ mag. This value agrees within the combined 1σ uncertainty with the true distance modulus derived from the Wesenheit index. Unless follow-up near-infrared data for the Cepheids in NGC 55 become available which will allow a more accurate determination of the mean reddening of the Cepheids due to the longer wavelength baseline, we will adopt, as our current best value from optical VI photometry, 26.40 mag as the true distance modulus of NGC 55.

5. Discussion

Our adopted distance to NGC 55 from optical photometry in the V and I bands of 130 long-period Cepheid variables discovered in the present survey is 26.40 mag, corresponding to 1.91 Mpc. This value is practically identical to the distance of NGC 300 of (1.88 ± 0.06) Mpc (Gieren et al. 2005a), strongly supporting the idea that NGC 300 and NGC 55 form a gravitationally bound system as previously suggested by Whiting (1999), and others. The intrinsic uncertainty of the distance modulus of NGC 55 derived from the (V-I) Wesenheit index is ± 0.05 mag, and is basically determined by the accuracy of our photometry. There are additional sources of systematic error which may affect this result. One of them is the effect of blending of some Cepheids in our sample with relatively bright stars, too close to the Cepheids to be resolved in our images. In a previous study of NGC 300 in which we could compare ground-based photometry of a large number of Cepheids to BVI photometry of the same variables obtained with the *Hubble Space Telescope* (Bresolin et al. 2005), we were able to show that the effect blending with unresolved bright stars has on the distance modulus derived from the ground-based images is less than 0.04 mag. We think that the effect of blending on our present distance result for NGC 55 should be of the same order given that the galaxy has the same distance as NGC 300, and the resolution of our NGC 55 images is very similar to the one of the images we used in our distance work for NGC 300. The higher inclination of NGC 55 with respect to the line of sight however might render the

effect due to blending somewhat more severe, as possibly indicated by the slightly enhanced dispersion of the V-band PL relation for NGC 55 (Fig. 4) as compared to NGC 300 (Gieren et al. 2004). We estimate that blending with unresolved bright companion stars might introduce a systematic uncertainty of ± 0.07 mag, in the case of NGC 55.

Another possible source of systematic uncertainty on our distance result is the slopes of the fiducial LMC Cepheid PL relations which we have adopted from the extensive work of the OGLE Project (Udalski 2000). Recent work by Ngeow and Kanbur (2006; see also references given therein) suggest that the LMC Cepheid PL relations in V and I are likely to show significant breaks at a period close to 10 days, leading to slightly different slopes of the PL relations valid for periods shorter, and longer than 10 days. The effect is very subtle and according to Ngeow and Kanbur (2006) affects distance moduli determined from the V and I PL relations by less than 0.03 mag. We will assume here that the systematic uncertainty of our NGC 55 distance modulus due to the uncertain slope of the fiducial PL relation in the LMC is ± 0.03 mag.

Another effect which might systematically affect our distance result is the possible metallicity dependence of the Cepheid PL relation. If such an effect is significant, the use of a PL relation calibrated for LMC metallicity for a Cepheid population whose metallicity is quite different from the LMC Cepheids would systematically affect the distance result. A number of recent theoretical and empirical studies have tried to trace down the effect of Cepheid metal abundances on their absolute magnitudes in different wavelength ranges (e.g. Storm et al. 2004; Saha et al. 2006), but unfortunately so far no clear-cut and accurate results have been obtained. A thorough discussion of the metallicity effect on the Cepheid PL relation has very recently been given by Mottini et al. (2006). Their own results obtained from relating the PL relation residuals for individual Cepheids to their metallicities measured from high-resolution spectra are consistent with a null effect of Cepheid iron abundance on the PL relation zero points in both the optical V and the near-infrared K band. It is clear, however, that further work on this very important question is needed which will eventually provide more accurate results than all previous work published over the past years. This is one of the principal goals of our Araucaria Project. What can be said at the present time is that from the results published over the past 8 years (see Mottini et al. 2006) it appears that the metallicity effect is small, not exceeding a few percent in any of the standard optical and near-IR photometric bands. This seems true both for the PL relation zero points (Pietrzynski and Gieren 2006), and for their slopes in different bands (e.g. Gieren et al. 2005c). We adopt ± 0.04 mag as an additional systematic uncertainty on our present distance modulus for NGC 55 from the metallicity effect.

Probably the largest systematic uncertainty on our present distance result for NGC 55

is the size of the reddening intrinsic NGC 55 which affects our Cepheid sample. The use of the Wesenheit index in this paper, and the consistency of the distance moduli from V, I and W using the reddening value derived from our data makes us expect that the effect of the uncertain intrinsic reddening on our adopted distance modulus is not larger than a few percent, however. It is clear that our planned follow-up distance determination for NGC 55 from accurate near-infrared photometry of a significant subsample of the Cepheids in our catalog will eliminate reddening as a significant source of error, as we could demonstrate in our previous Cepheid near-infrared studies in the Araucaria Project, and allow us to reduce the total systematic uncertainty on the distance of NGC 55 to a figure close to 3%. Moving our distance work to the near-infrared will also reduce the effect of blending because most stars which are bright enough to contaminate the fluxes of Cepheid variables at a significant level, including the secondary components in Cepheid binary systems, are expected to be blue stars which affect the Cepheid fluxes in optical bands more strongly than in the near-infrared (e.g. Gieren 1982; Szabados 1996).

Summarizing this discussion, we estimate that the total systematic uncertainty on our current distance modulus for NGC 55 from the various factors we have discussed is in the order of ± 0.09 mag. Our adopted true distance modulus to NGC 55 is therefore $26.40 \text{ mag} \pm 0.05 \text{ mag}$ (internal error) $\pm 0.09 \text{ mag}$ (systematic error). We note that this result is tied to an *assumed* distance modulus of 18.50 of the LMC and does not include the current uncertainty on the LMC distance, which is probably in the order of ± 0.10 mag as well. Since all the distances we are obtaining in the Araucaria Project are tied to the same assumed LMC distance of 18.50 mag, however, the *relative* distances between our target galaxies are not affected.

6. Conclusions

We have carried out an extensive search for Cepheid variables in NGC 55 which has resulted in the discovery of 143 bona fide Cepheids. We provide the periods of these objects and their mean magnitudes in the optical V and I bands, as well as their accurate positions. From 130 of these variables with periods between 10 and 100 days we obtain the period-luminosity relations in V and I, and for the reddening-independent (V-I) Wesenheit index. The distance moduli derived from these data demonstrate that apart from the very small foreground reddening, there is significant additional reddening which is produced inside NGC 55, which we determine as $E(B-V) = 0.102 \text{ mag}$ from our data. Our best adopted value for the distance modulus of NGC 55 from optical photometry of its large population of Cepheid variables is $26.40 \text{ mag} \pm 0.05 \text{ mag}$ (intrinsic) $\pm 0.09 \text{ mag}$

(systematic). The systematic uncertainty includes the estimated contributions from the effect of metallicity on the PL relation, the current uncertainty on the slope of the fiducial LMC Cepheid PL relation we have used in our distance determination, the effect of blending of the Cepheids in our sample with bright companion stars which are not resolved in our images, accuracy of the photometric zero points, and of dust absorption internal to NGC 55 affecting the optical Cepheid magnitudes. As in the previous papers of the Araucaria Project, our adopted distance modulus of our present target galaxy is tied to an assumed LMC true distance modulus of 18.50. Our estimated 0.09 mag systematic uncertainty of the distance modulus of NGC 55 does not contain the additional systematic uncertainty of the adopted value of the LMC distance.

The Cepheid distance derived in this paper for NGC 55 is more accurate than the other distance determinations for this nearby galaxy available in the literature which we discussed in the Introduction of this paper. In the near future, we expect to improve the accuracy of the distance determination to NGC 55 even further through near-IR photometry of a subsample of the Cepheids in our catalog, which will reduce the current uncertainties in our optical study due to reddening and blending.

Within the errors, the distance to NGC 55 from its Cepheid variables is *identical* to the Cepheid distance of 1.88 Mpc to NGC 300 (Gieren et al. 2005a) which is located close to NGC 55 in the sky. This strengthens the case for a physical association of NGC 55 and NGC 300, as previously suggested by a number of authors.

We are grateful to Las Campanas Observatory for the support provided to our observers, and to the CNTAC for providing the large amounts of telescope time which were necessary to complete this project. We also would like to thank the anonymous referee for interesting comments and suggestions. GP, WG, RM and AG gratefully acknowledge financial support for this work from the Chilean Center for Astrophysics FONDAP 15010003. WG also acknowledges support from the Centrum fuer Internationale Migration und Entwicklung in Frankfurt/Germany who donated the workstation on which a substantial part of the data reduction and analysis for this project was carried out. Support from the Polish KBN grant No 2P03D02123 and BST grant for Warsaw University Observatory is also acknowledged.

REFERENCES

- Alcaino, G. and Liller, W., 1984, AJ, 89, 814
 Bresolin, F., Pietrzyński, G., Gieren, W., and Kudritzki, R.P., 2005, ApJ, 634, 1020

- Gieren, W., 1982, *ApJS*, 49, 1
- Gieren, W., Pietrzyński, G., Walker, A., Bresolin, F., Minniti, D., Kudritzki, R.P., Udalski, A., Soszyński, I., Fouqué, P., Storm, J., and Bono, G., 2004, *AJ*, 128, 1167
- Gieren, W., Pietrzyński, G., Soszyński, I., Bresolin, F., Kudritzki, R.P., Minniti, D., and Storm, J., 2005a, *ApJ*, 628, 695
- Gieren, W., Pietrzyński, G., Bresolin, F., Kudritzki, R.P., Minniti, D., Urbaneja, M., Soszyński, I., Storm, J., Fouqué, P., Bono, G., Walker, A.R., and García, J., 2005b, *The Messenger*, 121, 23
- Gieren, W., Storm, J., Barnes, T.G., Fouqué, P., Pietrzyński, G., and Kienzle, F., 2005c, *ApJ*, 627, 224
- Gieren, W., Pietrzyński, G., Nalewajko, K., Soszyński, I., Bresolin, F., Kudritzki, R.P., Minniti, D., and Romanowsky, A., 2006, *ApJ*, 647, 1056
- Karachentsev, I.D., Grebel, E.K., Sharina, M.E., et al., 2003, *A&A*, 404, 93
- Mennickent, R.E., Gieren, W., Soszyński, I., and Pietrzyński, G., 2006, *A&A*, 450, 873
- Mottini, M., Primas, F., Romaniello, M., Bono, G., Groenewegen, M.A.T., and Francois, P., 2006, *A&A*, submitted
- Ngeow, C., Kanbur, S.M., 2006, *ApJ*, 642, L29
- Pietrzyński, G., Gieren, W., Fouqué, P., and Pont, F., 2002, *AJ*, 123, 789
- Pietrzyński, G., Gieren, W., and Udalski, A., 2002, *PASP*, 114, 298
- Pietrzyński, G., Gieren, W., Udalski, A., Bresolin, F., Kudritzki, R.P., Soszyński, I., Szymański, M., and Kubiak, M., 2004, *AJ*, 128, 2815
- Pietrzyński, G., Gieren, W., Soszyński, I., Bresolin, F., Kudritzki, R.P., Dall’Ora, M., Storm, J., and Bono, G., 2006a, *ApJ*, 642, 216
- Pietrzyński, G., Gieren, W., Udalski, A., Soszyński, I., Bresolin, F., Kudritzki, R.P., Mennickent, R.E., Kubiak, M., Szymański, M., and Hidalgo, S., 2006b, *ApJ*, in press
- Pietrzyński, G., and Gieren, W., 2006, *Mem. S.A. Italiana*, 75, 282
- Saha, A., Thim, F., Tammann, G.A., et al., 2006, *astro-ph/0602572*
- Schlegel, D.J., Finkbeiner, D.P., and Davis, M., 1998, *ApJ*, 500, 525
- Schwarzenberg-Czerny, A., 1989, *MNRAS*, 241, 153
- Soszyński, I., Gieren, W., Pietrzyński, G., Bresolin, F., Kudritzki, R.P., and Storm, J., 2006, *ApJ*, in press

- Storm, J., Carney, B.W., Gieren, W.P., Fouqué, P., Latham, D.W. and Fry, A.M., 2004, A&A, 415, 531
- Szabados, L., 1996, A&A, 311, 189
- Tuellmann, R., Rosa, M.R., Elwert, T., Bomans, D.J., Ferguson, A.M.N., and Dettmar, R.J., 2003, A&A, 412, 69
- Udalski, A., 2000, Acta Astron., 50, 279
- Udalski, A., 2003, Acta Astron., 53, 291
- Udalski, A., Soszyński, I., Szymański, M., Kubiak, M., Pietrzyński, G., Woźniak, P., and Żebruń, K., 1999, Acta Astron., 49, 223
- Urbaneja, M.A., Herrero, A., Bresolin, F., Kudritzki, R.-P., Gieren, W., Puls, J., Przybilla, N., Najarro, F. and Pietrzyński, G., 2005, ApJ, 622, 862
- Van de Steene, G.C., Jacoby, G.H., Praet, C., Ciardullo, R. and Dejonghe, H., 2006, A&A, in press
- Whiting, A.B., 1999, AJ, 117, 202
- Woźniak, P., 2000, Acta Astron., 50, 421




Fig. 1.— The location of the observed field in NGC 55 on the DSS blue plate. The field of view was about 35×35 arcmin. North is up and East is to the left. The locations of the Cepheids discovered in our survey are indicated by the white circles.

Table 1. Comparison of our V band photometry

ID	V [mag] Alcaino and Liller	V [mag] this paper	Remarks
V	16.05	16.054	Variable star
W	16.20	16.212	
X	16.95	17.033	
T	15.79	15.796	
K	14.47	14.448	
N	14.90	14.919	
O	14.91	14.916	

Table 2. Cepheids in NGC 55

ID	RA (J2000)	DEC (J2000)	P [days]	T ₀	< V > [mag]	< I > [mag]	< W _I > [mag]
cep001	0:14:12.95	-39:08:41.7	175.9086	2452808.50787	19.249	18.409	17.107
cep002	0:15:12.00	-39:12:18.0	152.0943	2452746.27372	19.562	18.613	17.142
cep003	0:14:36.57	-39:11:08.9	112.6964	2452800.19911	20.177	19.131	17.510
cep004	0:15:14.28	-39:13:17.4	97.7291	2452823.21391	20.536	19.288	17.354
cep005	0:15:10.11	-39:12:26.0	85.0550	2452769.04780	20.835	19.456	17.319
cep006	0:15:47.32	-39:16:10.7	76.6842	2452797.49527	20.101	19.147	17.668
cep007	0:15:40.34	-39:15:08.0	75.8328	2452788.65035	20.336	19.510	18.230
cep008	0:14:45.35	-39:13:17.4	74.6393	2452828.22484	20.124	19.253	17.903
cep009	0:15:13.99	-39:12:33.4	73.5323	2452782.28141	20.912	19.706	17.837
cep010	0:14:06.59	-39:08:11.6	66.8528	2452788.45946	20.356	19.464	18.081
cep011	0:13:58.27	-39:08:22.8	62.7910	2452799.95331	20.432	19.568	18.229
cep012	0:15:14.93	-39:12:52.5	62.3186	2452811.59034	20.834	19.665	17.853
cep013	0:14:12.72	-39:09:27.2	57.1544	2452802.83374	20.425	19.297	17.549
cep014	0:15:56.86	-39:15:23.6	57.0377	2452840.41529	20.228	19.252	17.739
cep015	0:14:54.55	-39:12:49.4	56.9074	2452832.44085	20.418	19.550	18.205
cep016	0:14:37.74	-39:10:55.9	53.1177	2452834.23720	20.746	19.595	17.811
cep017	0:15:00.36	-39:12:25.9	48.7260	2452809.41765	20.910	19.877	18.276
cep018	0:16:04.10	-39:16:50.6	46.3493	2452807.55483	21.349	20.094	18.149
cep019	0:15:48.48	-39:15:41.1	46.2082	2452838.40323	20.506	19.741	18.555
cep020	0:14:25.63	-39:09:57.3	44.0077	2452838.42241	21.268	20.181	18.496
cep021	0:15:41.65	-39:15:15.1	42.7678	2452808.11211	20.987	19.964	18.378
cep022	0:15:00.42	-39:12:26.5	41.7433	2452829.14536	21.095	20.401	19.325
cep023	0:14:18.63	-39:10:39.3	41.7395	2452811.35926	21.122	20.469	19.457
cep024	0:14:37.93	-39:10:52.2	41.4594	2452818.60556	20.952	20.057	18.670
cep025	0:15:04.49	-39:12:37.4	40.2691	2452829.58075	20.561	19.784	18.580
cep026	0:14:43.45	-39:11:14.8	39.0861	2452837.83586	21.419	20.469	18.996
cep027	0:14:39.76	-39:10:41.7	38.7170	2452814.19686	21.456	20.152	18.131
cep028	0:14:38.17	-39:11:30.5	37.8170	2452817.09891	21.316	20.365	18.891
cep029	0:14:55.46	-39:13:18.8	36.3548	2452818.79637	20.800	20.161	19.171
cep030	0:15:46.52	-39:16:18.3	36.1691	2452822.55861	21.621	20.572	18.946
cep031	0:14:42.58	-39:11:28.4	35.6486	2452842.23736	20.970	20.641	20.131
cep032	0:14:42.52	-39:11:27.6	35.5466	2452820.18259	19.621	18.851	17.657
cep033	0:16:04.21	-39:16:30.6	35.0516	2452848.08098	21.889	20.207	17.600

Table 2. Cepheids in NGC 55 - continued

ID	RA (J2000)	DEC (J2000)	P [days]	T ₀	< V > [mag]	< I > [mag]	< W _I > [mag]
cep034	0:14:44.36	-39:11:21.5	34.5518	2452816.49099	20.663	20.151	19.357
cep035	0:14:53.98	-39:12:09.2	34.1178	2452825.84880	99.999	20.277	99.999
cep036	0:15:11.80	-39:12:52.4	32.8210	2452821.22415	21.134	20.064	18.405
cep037	0:14:25.31	-39:10:14.9	32.4267	2452827.27556	20.915	20.026	18.648
cep038	0:14:59.73	-39:11:54.7	31.5292	2452822.25693	21.746	20.608	18.844
cep039	0:14:19.43	-39:09:46.7	31.1574	2452850.00292	21.076	20.395	19.339
cep040	0:15:02.95	-39:12:27.5	31.0364	2452845.00824	21.143	20.337	19.088
cep041	0:14:58.86	-39:12:50.0	30.2467	2452826.35274	21.029	20.335	19.259
cep042	0:15:33.90	-39:14:48.5	29.5002	2452830.30335	20.976	20.290	19.227
cep043	0:16:02.28	-39:16:18.2	29.4512	2452837.34388	21.908	20.578	18.516
cep044	0:15:54.86	-39:15:19.3	28.0657	2452848.53476	21.433	20.510	19.079
cep045	0:14:54.02	-39:12:19.3	28.0509	2452841.89663	21.048	20.446	19.513
cep046	0:14:38.92	-39:11:05.1	27.7661	2452822.84443	21.288	20.349	18.894
cep047	0:14:42.78	-39:11:26.6	27.2861	2452830.27499	20.623	20.283	19.756
cep048	0:14:50.55	-39:12:22.7	27.1599	2452831.66545	21.084	20.283	19.041
cep049	0:15:40.28	-39:13:39.8	26.6976	2452838.50199	21.564	20.735	19.450
cep050	0:15:34.45	-39:15:13.7	26.3878	2452826.21631	21.234	20.605	19.630
cep051	0:15:37.86	-39:14:48.6	26.3199	2452826.63416	22.733	21.131	18.648
cep052	0:15:30.94	-39:12:34.0	26.0379	2452849.86527	21.643	20.511	18.756
cep053	0:14:46.23	-39:11:43.5	25.5505	2452845.03937	21.430	20.503	19.066
cep054	0:15:02.18	-39:14:04.5	25.4368	2452834.77264	21.014	20.531	19.782
cep055	0:15:16.57	-39:12:46.6	25.1401	2452840.13751	21.761	20.598	18.795
cep056	0:15:59.28	-39:16:12.6	24.9857	2452833.12543	21.693	20.621	18.959
cep057	0:16:04.32	-39:16:46.6	24.5666	2452840.29535	22.032	20.648	18.503
cep058	0:15:30.03	-39:12:20.3	23.9016	2452835.76728	22.219	20.912	18.886
cep059	0:15:53.41	-39:16:08.4	23.7508	2452832.09632	22.276	20.976	18.961
cep060	0:14:44.84	-39:11:08.7	23.7096	2452829.85330	22.218	21.210	19.648
cep061	0:15:52.16	-39:16:37.4	23.5439	2452826.41263	21.592	20.677	19.259
cep062	0:15:33.54	-39:13:28.7	23.5351	2452841.77805	21.755	20.371	18.226
cep063	0:14:42.37	-39:10:46.0	23.3327	2452846.32745	21.163	20.695	19.970
cep064	0:15:55.80	-39:16:35.7	22.9247	2452843.72260	21.896	20.719	18.895
cep065	0:14:46.56	-39:11:12.9	22.8332	2452846.49672	22.304	20.908	18.744
cep066	0:15:32.71	-39:14:39.4	22.6913	2452845.71810	20.965	20.145	18.874
cep067	0:14:34.49	-39:10:25.3	22.4043	2452836.82067	21.495	20.749	19.593

Table 2. Cepheids in NGC 55 - continued

ID	RA (J2000)	DEC (J2000)	P [days]	T ₀	< V > [mag]	< I > [mag]	< W _I > [mag]
cep068	0:14:49.80	-39:13:23.1	22.2934	2452844.54875	22.006	21.177	19.892
cep069	0:14:25.07	-39:09:10.8	22.2824	2452846.57642	21.753	99.999	99.999
cep070	0:15:10.89	-39:11:59.2	22.1505	2452847.40035	22.468	21.136	19.071
cep071	0:15:38.89	-39:14:56.5	22.0950	2452833.70839	21.252	20.785	20.061
cep072	0:15:18.53	-39:12:29.6	21.9627	2452837.12408	99.999	21.118	99.999
cep073	0:14:36.52	-39:12:19.1	21.8533	2452832.62534	21.937	21.036	19.639
cep074	0:15:50.28	-39:15:46.1	21.5309	2452831.85886	21.263	21.303	21.365
cep075	0:14:45.91	-39:11:27.6	21.2743	2452842.58589	22.084	20.868	18.983
cep076	0:16:07.66	-39:17:28.2	21.2415	2452846.95235	22.539	20.933	18.444
cep077	0:14:02.69	-39:11:09.3	20.5319	2452831.18683	21.189	20.648	19.809
cep078	0:15:44.34	-39:15:12.2	20.3943	2452835.39236	21.673	20.826	19.513
cep079	0:14:17.12	-39:11:41.9	20.3942	2452845.84048	20.952	20.166	18.948
cep080	0:14:44.24	-39:12:45.4	20.2633	2452845.15643	21.504	20.859	19.859
cep081	0:15:18.26	-39:12:14.1	19.8080	2452842.48408	22.407	21.007	18.837
cep082	0:15:49.29	-39:15:48.3	19.3776	2452838.69568	21.852	20.947	19.544
cep083	0:14:58.69	-39:11:54.3	19.3580	2452847.98353	21.926	99.999	99.999
cep084	0:14:55.31	-39:12:50.8	19.3436	2452836.30750	21.131	21.072	20.981
cep085	0:14:46.96	-39:12:37.3	19.0477	2452834.05731	21.426	20.924	20.146
cep086	0:15:53.91	-39:16:25.3	18.5752	2452836.59181	21.745	20.478	18.514
cep087	0:14:20.17	-39:11:24.2	18.3966	2452847.63746	21.764	21.053	19.951
cep088	0:15:16.02	-39:11:53.3	18.0795	2452834.01755	22.852	21.235	18.729
cep089	0:14:46.51	-39:10:42.0	17.9405	2452835.18692	21.352	20.453	19.060
cep090	0:14:40.23	-39:10:03.3	17.8206	2452847.60431	21.811	20.837	19.327
cep091	0:14:59.60	-39:11:48.9	17.6201	2452847.29662	99.999	21.131	99.999
cep092	0:14:29.65	-39:10:16.4	17.5669	2452837.70808	22.679	21.178	18.851
cep093	0:15:29.94	-39:15:06.4	17.5499	2452833.43163	21.819	20.561	18.611
cep094	0:15:31.13	-39:15:40.7	17.2671	2452847.02598	21.770	20.835	19.386
cep095	0:14:48.08	-39:11:25.8	17.2338	2452842.19519	21.809	20.881	19.443
cep096	0:15:22.52	-39:13:31.9	17.1511	2452833.75601	21.982	21.323	20.302
cep097	0:14:46.82	-39:10:51.6	17.0303	2452839.29756	21.417	20.918	20.145
cep098	0:14:14.50	-39:10:42.0	16.9912	2452844.98131	21.759	20.945	19.683
cep099	0:14:27.99	-39:11:46.5	16.9451	2452849.39100	22.094	21.401	20.327
cep100	0:14:20.95	-39:10:52.3	16.5556	2452836.29047	21.822	21.572	21.184
cep101	0:14:22.12	-39:10:50.0	16.5403	2452848.50372	21.261	20.851	20.215

Table 2. Cepheids in NGC 55 - continued

ID	RA (J2000)	DEC (J2000)	P [days]	T ₀	< V > [mag]	< I > [mag]	< W _I > [mag]
cep102	0:14:49.55	-39:12:20.4	15.9295	2452834.61847	21.287	20.208	18.536
cep103	0:16:00.34	-39:16:13.5	15.8413	2452840.15070	21.558	20.960	20.033
cep104	0:15:48.09	-39:15:48.6	15.4922	2452848.08276	22.004	20.891	19.166
cep105	0:15:39.21	-39:13:23.9	15.3373	2452839.24376	22.385	21.340	19.720
cep106	0:15:50.88	-39:17:01.9	15.0940	2452840.27171	22.547	21.535	19.966
cep107	0:14:26.94	-39:10:37.8	15.0797	2452837.95823	22.215	21.460	20.290
cep108	0:15:15.19	-39:12:13.9	14.9736	2452847.98548	22.415	21.237	19.411
cep109	0:14:48.72	-39:13:33.0	14.9476	2452838.39412	21.990	21.291	20.208
cep110	0:15:24.97	-39:12:51.2	14.6266	2452852.25159	99.999	20.595	99.999
cep111	0:14:33.45	-39:12:33.2	14.3815	2452844.42839	22.188	21.052	19.291
cep112	0:14:34.73	-39:12:26.8	14.1692	2452847.42774	22.431	21.514	20.093
cep113	0:15:58.40	-39:16:12.1	14.1150	2452849.96568	22.668	20.804	17.915
cep114	0:15:49.00	-39:16:04.5	14.0094	2452846.51687	22.405	21.591	20.329
cep115	0:14:30.72	-39:09:49.6	13.9050	2452848.72358	22.154	21.039	19.311
cep116	0:15:16.28	-39:13:27.1	13.5937	2452850.69133	22.168	21.956	21.627
cep117	0:15:25.45	-39:15:04.2	13.4465	2452840.17843	22.937	21.299	18.760
cep118	0:14:28.15	-39:10:10.0	13.2328	2452849.85684	21.867	20.964	19.564
cep119	0:15:46.71	-39:16:35.1	13.0498	2452847.75940	22.813	99.999	99.999
cep120	0:14:27.44	-39:12:09.3	13.0361	2452849.11512	21.542	21.077	20.356
cep121	0:14:08.26	-39:12:23.1	12.4816	2452849.92789	22.363	21.678	20.616
cep122	0:16:04.38	-39:16:19.6	12.4326	2452850.09452	22.454	21.499	20.019
cep123	0:15:41.26	-39:14:57.7	12.3569	2452848.82971	21.817	21.271	20.425
cep124	0:14:57.99	-39:12:37.2	12.0533	2452851.65153	21.621	21.122	20.349
cep125	0:14:42.00	-39:12:31.5	11.9959	2452851.18123	21.933	21.423	20.632
cep126	0:14:16.76	-39:09:09.9	11.9130	2452840.80674	22.553	21.662	20.281
cep127	0:14:34.94	-39:12:12.3	11.4633	2452842.70761	22.121	21.197	19.765
cep128	0:16:11.77	-39:16:17.3	11.0061	2452849.74823	22.051	21.401	20.393
cep129	0:15:57.70	-39:16:49.4	10.8421	2452852.89578	22.755	21.686	20.029
cep130	0:16:13.34	-39:16:49.0	10.7971	2452842.40920	22.413	21.585	20.302
cep131	0:14:19.89	-39:11:16.4	10.6378	2452841.66009	22.031	21.115	19.695
cep132	0:16:06.00	-39:15:46.3	10.2913	2452846.29574	22.733	21.809	20.377
cep133	0:15:14.60	-39:13:20.3	10.2782	2452852.15148	22.032	20.869	19.066

Table 2. Cepheids in NGC 55 - continued

ID	RA (J2000)	DEC (J2000)	P [days]	T ₀	< V > [mag]	< I > [mag]	< W _I > [mag]
cep134	0:14:29.79	-39:12:24.6	9.1664	2452848.79550	22.448	21.789	20.768
cep135	0:14:51.96	-39:11:44.5	8.8605	2452853.86214	19.997	19.931	19.829
cep136	0:13:55.94	-39:10:57.1	8.8481	2452847.89737	22.383	22.206	21.932
cep137	0:16:08.89	-39:15:26.8	8.2505	2452841.86954	22.710	21.937	20.739
cep138	0:14:19.33	-39:10:36.4	7.9824	2452843.03592	22.466	21.714	20.548
cep139	0:16:25.33	-39:16:27.7	7.7652	2452848.40174	22.798	22.382	21.737
cep140	0:14:35.99	-39:12:22.9	7.3029	2452846.30757	22.303	21.148	19.358
cep141	0:15:12.74	-39:14:54.5	6.6286	2452845.48814	99.999	22.250	99.999
cep142	0:16:00.27	-39:14:59.4	6.1346	2452846.62733	22.869	22.188	21.132
cep143	0:14:40.81	-39:11:35.3	5.5692	2452848.85807	22.832	21.525	19.499

Table 3. Individual V and I Observations

object	filter	HJD	mag	σ_{mag}
cep001	V	2452850.80482	19.242	0.008
cep001	V	2452853.83449	19.249	0.007
cep001	V	2452858.82639	19.283	0.008
cep001	V	2452860.77519	19.261	0.013
cep001	V	2452861.92803	19.277	0.012
cep001	V	2452868.74762	19.320	0.012
cep001	V	2452872.73327	19.319	0.008
cep001	V	2452880.74169	19.374	0.009
cep001	V	2452886.72905	19.415	0.013
cep001	V	2452904.67335	19.449	0.010
cep001	V	2452912.67723	19.439	0.010
cep001	V	2452917.63772	19.448	0.015
cep001	V	2452921.63502	19.437	0.027
cep001	V	2452923.65303	19.429	0.021
cep001	V	2452931.63660	19.420	0.010
cep001	V	2452935.60175	19.396	0.011
cep001	V	2452944.69719	19.341	0.014
cep001	V	2452951.55163	19.259	0.021
cep001	V	2452956.54067	19.233	0.008
cep001	V	2452970.52595	19.104	0.009

Note. — The complete version of this table is in the electronic edition of the Journal. The printed edition contains only the the first 20 measurements in the V band for the Cepheid variable cep001.




Fig. 2.— Phased V- and I-band light curves for some Cepheids of different periods in our NGC 55 catalog. These light curves are representative for the light curves of other Cepheid variables of similar periods.




Fig. 3.— The V,V-I color-magnitude diagram for NGC 55. The Cepheids discovered in our survey are marked with filled circles.




Fig. 4.— The period-luminosity relation for NGC 55 Cepheids in the V band. The slope of the relation has been adopted from the LMC Cepheids (see text). We have used the 130 Cepheids with periods between 10 and 100 days for the distance determination to NGC 55.




Fig. 5.— Same as Fig. 4, for the I band.




Fig. 6.— Same as Fig. 4, for the reddening-independent (V-I) Wesenheit magnitudes.